# Herschel-PACS observation of gas lines from the disc around HD141569A

## Conference or Workshop Item

# Herschel-PACS observation of gas lines from the disc around HD141569A

**W.-F. Thi**[1], C. Pinte[1], E. Pantin[2], J.C. Augereau[1], G. Meeus[3], F. Ménard[1,4], C. Martin-Zaïdi[1], P. Woitke[5], P. Riviere-Marichalar[6] I. Kamp[6], A. Carmona[1], G. Sandell[7], C. Eiroa[3], W. Dent[8], B. Montesinos[3], G. Aresu[6], R. Meijerink[6] , M. Spaans[16] , G. White[9,10], D. Ardila[11] , J. Lebreton[1], I. Mendigutía[12], S. Brittain[12]

1 UJF-Grenoble 1 / CNRS-INSU, Institut de Planétologie et d'Astrophysique (IPAG) UMR 5274, Grenoble, F-38041, France
e-mail: Wing-Fai.Thi at obs.ujf-grenoble.fr
[2] CEA, Saclay, Fr; [3] UAM, Spain; [4] UMI-LFCA,CNRS/INSU, Chile; [5] SUPA, University of St-Andrews, UK; [6] Kapteyn Astronomical Institute, NL; [7] SOFIA-USRA, NASA, USA; [8] ALMA, Chile; [9] Open University, UK; [10] RAL, UK; [11] NASA Herschel Center, USA, [12] Clemson University, USA

Protostars and Protoplanets VI, 15-20 July 2013 Heidelberg

HD 141569A was observed by Herschel as part of the Gas in Protoplanetary Discs Survey (Dent et al. 2013). We complemented with ground-based observations to constrain the gas and dust in the disc.

## Herschel-PACS GASPS programme : [OI] 63, 145 & [CII] lines detected




## Model parameters

B9.5V star, 5 Myrs, d=108pc
$M_{dust}$(disc) ~ 2.6 x $10^{-6}$ $M_{Sun}$
$M_{gas}$(disc) ~2.5 x $10^{-4}$ $M_{Sun}$
$M_{PAH}$(disc) ~ 1.8 x $10^{-12}$ $M_{sun}$
Inner disc: 5-110 AU
Outer disc: 185-500 AU

Gas mass/Dust mass ~ 100

## Density structure consistent with VISIR image at 8.6 micron




PAH emission at 8.6 micron

## Disc continuum modelling with MCFOST

Fit to the SED + PAH features with MCFOST (Pinte et al. 2006)

nlte PAH + dust opacities treated simultaneously

PAH treatment: Draine & Li




Cut along the North-South direction: the PAH emission is resolved




Inner disc outer edge at 110 AU

## Gas chemistry and line transfer modelling with ProDiMo

[OI] 63, [OI] 145, [CII], and CO 3-2: flat, gas-to-dust~100 disc models produce fluxes within a factor 2 except for [OI] 63.




Comparison modelled CO 3-2 profile with observations (Dent et al. 2005)




Reference:
HD141569, article submitted to A&A
MCFOST: Pinte et al. 2006 A&A 459, 797
ProDiMo: Wotike et al. 2009 A&A 501, 383
Dent et al. 2005 2005, MNRAS, 359, 663
GASPS project: Dent, Thi, Kamp, et al., 2013, PASP, 125, 477
DIANA project: http://www.diana-project.com/

## Conclusions

- From the PAH image, the inner disc extents to at least 110 AU.
- All models with gas-to-dust mass ratio from 10 to100 overpredict the [OI] 63 micron flux. The oxygen chemistry may need to be revised.
- A model with gas-to-dust mass ratio of 100 is consistent with all the other gas constraints.
- Disc models with low opening angles (H/r) are favored due to the sensitity of the [CII] and CO 3-2 flux on the gas density (flt discs are denser).

Acknowledgment: E. van Dishoeck for discussion, European contract EU FP7-2011 Grant Agreement nr. 284405 (PERG06-GA-2009-256513), ANR contracts ANR-07-BLAN-0221, ANR-2010-JCJ-0504-01, ANR CHEX2011 SEED, Millenuim Science Initiative "Nucleus P10-022-F", Spanish grant AYA 2011-26202. Calculations were done on Fostino funded by ANR.

Please contact me for more details. I am also looking for a tenure/tenure-track position.